



Department of Horticulture

Purdue University Cooperative Extension Service • West Lafayette, IN

Chinese Vegetables

*Michael N. Dana**

Foreign cuisines are of great interest to the American cook, but high quality, authentic fresh ingredients are sometimes difficult to find. Authentic Chinese cooking demands fresh vegetables which may only be available to you if you grow them yourself. Yet their use is not limited to Chinese cuisine. Many can be used as substitutes in or tasty additions to traditional American dishes.


The culture of most Chinese vegetables is not complicated. Chinese types of beans, squash, cucurbits (cucumber), and cole crops (broccoli, cabbage) are closely related to their "Western" counterparts and have similar cultural requirements (Table 1).

Chinese Water Garden

The Chinese water garden is an exception to the foregoing rule. Chinese lotus, water chestnuts, violet

stemmed taro, and arrowhead require a wet growing environment, contrary to the conditions commonplace in American vegetable gardens. In the Orient, these aquatic plants are grown in flooded rice paddies.

You can simulate the conditions that these plants require with a little innovation. A typical approach is to use a 25-gallon minimum capacity container for water in which smaller plant containers are submerged (except for lotus). The soil should be typical garden soil, not a peat-containing potting mix. You will also need to include some submerged oxygenating plants (SOP) and goldfish. The latter will help keep the water clear and insect free, and will be attractive in the water garden. SOP provide extra oxygen, which otherwise might be a limiting factor in a water environment. Prop the smaller plant containers up on bricks or rocks so the soil line is the proper depth below the water surface. Table 2 gives detailed cultural information on each crop.



** This publication was originally co-authored by James Simon.*

Table 1. Chinese Vegetables Suitable for the Conventional American Vegetable Garden.

English Name/ Chinese Name/ Latin Name	Seed Depth (inches)	Distance Between Plants/ Between Rows (inches)	Planting Dates South (Spring range/ Fall range)	Planting Dates North (Spring range/ Fall range)	Days to Maturity	Comments/Use
Azuki Bean Hong dow <i>Vigna angulans</i>	1/2-1	2-3/18-30	4-1/6-30 7-1/8-1	5-15/6-15 6-15/6-30	120	May be eaten as sprouts, fresh pickled or dried. Edible pods. Beans have slightly sweet flavor. Usually prepared as paste for baking. Highly nutritious (25% protein).
Amaranth Cien choy <i>Amaranthus tricolor</i>	1/4	10-18/18	2-15/4-1 8-20/10-10	3-20/4-20 9-1/10-1	30-90	Brilliantly colored leaves usually grown as an ornamental. Use as spinach, raw or cooked. Also called tampala.
Asparagus Pea Bin dow <i>Psophocarpus tetragonolobus</i>	1/2	6/18	2-20/3-20 --	3-20/5-1 8-1/9-15	50	Tastes like asparagus. Harvest pods when 1 in. long. Use in stir fry and soups. Whole plant, including flowers, edible. Also called goa bean, princess pea, winged bean.
Bitter Melon Koo gwa <i>Momordica charantia</i>	1	36-48/48-60	4-20/6-1 8-15/10-1	5-15/6-15 7-15/9-1	60-75	Ornamental vine that produces bitter quinine containing fruit that must be parboiled to eat. Use in stir fry and soups. Also called balsam pear, la-kwa, and bitter gourd. Extracts of plant have been used for medicinal purposes.
Broccoli, Chinese Gai lan <i>Brassica oleracea</i> Alboglabra Group	1/2	6/12	3-15/4-5 7-1/8-15	4-1/5-1 7-1/8-1	60-80	Stalks split and prepared as common broccoli. If peeled and split, stems cook faster. Also called Chinese kale.
Burdock Ngan pong <i>Arctium lappa</i>	1/2-1	6/20	3-1/4-1 8-1/9-1	3-15/4-10 7-1/7-20	45	Although roots generally eaten, the young leaves and stems are edible. Scrub or scrape to prevent darkening and remove bitterness. Crisp texture remains even in cooked dishes.
Celery Cabbage, Chinese Pai tsai <i>Brassica rapa</i> Pekinensis Group	1/2-1	14-30/30-36	3-1/4-1 8-1/9-1	3-15/4-10 7-1/7-20	70-80	Use as regular heading cabbage stir fried or pickled. Forms tight upright head.
Chives, Chinese, Garlic Gow choy <i>Allium tuberosum</i>	1/2	8/12-14	3-15/4-5 --	4-1/5-1 --	24-90	Stronger flavor than ordinary chives. Decorative plant and flowers. Also called oriental garlic.

Cucumbers, Chinese Tseng gwa <i>Cucumis sativus</i>	1/2-3/4	36/48	4-20/6-1 6-1/8-1	5-15/6-15 6-1/7-15	60	Small seeded fruit is original "burpleess" type. Eat raw, pickled or cooked. Mature fruits are stuffed. Young fruits with flowers attached are eaten as appetizers.
Eggplant, Chinese Ai gwa <i>Solanum melongena</i> var. <i>esculentum</i>	1/2	18-24/30-36	5-1/6-1 6-1/7-1	5-15/6-10 6-1/7-1	60-75	Small white fruit on some varieties. Good for pickling, sauteeing, or frying.
Fava Beans Fu dow <i>Vicia faba</i>	2 1/2	4-6/18-24	4-1/6-30 7-1/8-1	5-15/6-15 6-16/6-30	65-90	Young pods edible, may be shelled like peas or left to produce dry beans.
Flowering Cabbage Hwa choy <i>Brassica oleracea</i> Acephala Group	1/2	8-10/6	3-1/4-1 8-1/9-1	3-15/4-10 7-1/7-20	55-75	Shred, boil, bake, stuff as regular cabbage. Beautiful red, pink, and green combinations of leaves often grown as ornamentals.
Fuzzy Gourd Jiet gwa <i>Benincasa hispida</i>	1	36/48	4-20/6-1 6-1/8-1	5-16/6-15 6-1/7-15	75-85	Used in ways similar to zucchini or squash. Fruit must have fuzz and skin removed before being eaten. Also called Chinese watermelon, Chinese preserving melon.
Garland Chrysanthemum Tong how tsai <i>Chrysanthemum coronarium</i>	1/4-1/2	6-12/18	2-20/3-20 8-15/10-1	3-20/5-1 7-15/9-1	25-60	Leaves and stems are used as greens for a distinctive flavor. Dried flower petals are used in soups and stir fried dishes.
Giant Garlic Suan <i>Allium sativum</i>	2	12/12	2-20/3-20 --	3-15/4-15 --	180	Milder flavor than common garlic. May be eaten raw or blended into butter. Shoots may be used like chives.
Ginger Giang <i>Zingiber officinale</i>	3	16/24	5-1/6-1 --	5-15/6-10 --	90-130	Rhizome used in all types of dishes from meats to confections. Added to stir-fry oil before vegetables, it imparts a "warm" flavor.
Hot Peppers La chiao <i>Capiscum frutescens</i>	1/4	18-24/24-36	5-1/6-1 6-1/7-20	5-15/6-10 6-1/7-10	65-85	Use with discretion as a seasoning. Very hot fruit is used raw or dried.
Multiplier Onions Chang fa <i>Allium fistulosum</i>	1/2	10/18	3-1/4-1 --	3-10/4-10 --	55-120	Use as scallions in stir fry or raw on relish tray. Mild onion flavor. Hollow leaves may reach 3 ft.
Mung Beans Nga choy <i>Vigna radiata</i>	1/2	6-8/24	4-1/6-30 7-1/8-1	5-15/6-15 6-15/6-30	90-120	May be eaten as sprouts (Yah tsai) or in a later stage such as edible green pods. Beans will be sticky if boiled.

Mustard, Chinese	1/4	4-10/12	3-15/5-15	4-1/6-1	35-50	Use as salad green. Commonly pickled, but may be stir-fried.
Gai choy			8-15/10-1	7-15/9-1		
<i>Brassica juncea</i>						
Mustard Cabbage,	1/4	6/18	3-1/4-1	3-15/4-10	35-50	Leaves cooked as spinach, stalks used like asparagus. Growth habit similar to celery or chard.
Chinese Bok choy			8-1/9-15	7-15/8-15		
<i>Brassica rapa</i>						
Chinensis Group						
Okra, Chinese (Luffa)	1	36-48/48-60	4-20/6-1	5-15/6-15	115	Prepare immature fruit as squash or cucumber. Pare off ridges, but leave other skin intact. Mature fruit can be dried and used as sponges. Highly prolific, vigorous vine.
Shuh gwa			6-1/8-1	6-1/7-15		
<i>Luffa acutangula</i>						
Parsley, Chinese	1/4	4-6/18	3-10/4-10	4-1/5-1	60	Seeds have citrus-spice flavor, Add leaves as a seasoning. Plant is also called coriander. Chopped leaves especially fine with poached fish.
Yfen tsai			8-1/9-15	7-15/8-15		
<i>Coriandrum sativum</i>						
Pickling Melon	1/2	36-48/60-120	4-20/6-1	5-15/6-15	65	Use in any standard pickling recipe. Can also be cooked or eaten raw as a vegetable.
Chung choy			6-1/8-1	6-1/7-15		
<i>Cucumis melo</i>						
Conomon Group						
Pol herb Mustard, Chinese	1/4	12/18	3-15/5-15	4-1/6-1	30-60	Leafy foliage attractive in salads. Rich in vitamins.
Mizuna (Japanese name)			8-15/10-1	7-15/9-1		
<i>Brassica juncea</i> var. <i>crispifolia</i>						
Pumpkins, Chinese	1	18-36/60-90	4-20/6-1	5-1/5-30	130	Used in some ways as squash and pumpkins in soups, pies, baked or fried. Can be stored all winter.
Nang gwa			6-1/8-1	6-1/7-15		
<i>Cucurbita pepo</i>						
Radish, Chinese	1	2-10/12-30	3-1/5-1	3-20/5-10	60-80	Grate, steam, stir fry, pickle, boil or sculpture fleshy root. Leafy tops edible also. Large and very long root. Also called Japanese daikon. Many cultivars are available.
Luo buo			3-15/10-15	8-1/10-1		
<i>Raphanus sativus</i> 'Longipinnatus'						
Sesame	1/2	9/36	4-20/6-1	5-15/6-15	30-45	Seeds eaten raw or roasted. Oil from seed has nutty flavor and is the oil of stir-frying or deep frying.
Chin ma			7-1/7-15	6-15/7-20		
<i>Sesamum indicum</i>						
Snow Peas	2	2-3/18-30	2-20/3-20	3-20/5-1	50-85	Use as soon as picked. Combines well with fish dishes. Add to stir-fry only in last few seconds of cooking.
Ho lan dow			--	--		
<i>Pisum sativum</i> var. <i>macrocarpon</i>						
Soybeans	1/2	2-3/24-30	5-1/6-30	5-15/6-15	75-115	Mildly nutty flavor. May be served as sprouts, roasted or dried. Harvest green or dry.
Huang dow			6-1/7-15	6-1/7-5		
<i>Glycine max</i>						

Sweet Melons	1	18-36/60-72	4-20/6-1	5-15/6-15	115-130	Similar to cantaloupe or honeydew.
Chung Gwa <i>Cucumis melo</i>			7-17-15	6-15/7-20		
Winter Melon Doong gwa <i>Benincasa hispida</i>	1	18-36/48-60	4-20/6-1	5-15/6-15	150	All parts (young leaves, flower buds, both immature and mature fruit) may be used.
Yard-Long Beans Gauk dow <i>Vigna unguiculata</i> spp. <i>sesquipedalis</i>	1/2-1	48-60/24	4-20/6-30	5-10/6-30	60-90	12 to 16-inch beans can be used as common green beans. Grow on trellis or teepee supports to 8 ft. high.

Table 2. Chinese Water Garden Vegetables.

English Name/ Chinese Name/ Latin Name/ Pot Size (inches)	Minimum Pot Size (inches)	Depth to Plant in Soil (inches)	Water Depth Over Soil (inches)	When to Plant	What to Plant	What/When to Harvest	Comments/Use
Arrowhead Chee koo <i>Sagittaria sagittifolia</i>	7 diam 5 deep	1	3-6	after danger of frost	tuber	tuber/fall year	Tubers are not edible raw. Like a slightly nutty sweet potato, eat roasted or boiled. Also called swamp potato.
Chinese Lotus Lien ou <i>Nelumbo nucifera</i>	25 gal	2 (with growing tip visible on soil surface)	10	after danger of frost	rhizome or by seeds	buds, flowers, seed pods/as soon as formed/ rhizomes (or root)	All plant parts can be used, raw or cooked, as greens, in soups, etc. The "roots" (swollen stems which grow in the mud) are most often used in fried, baked, or raw form. They resemble huge sausage links, reddish brown in color.
Violet Stemmed Taro Yu tou <i>Xanthosoma violaceum</i>	7 diam 5 deep	1	3-6	after danger of frost	tuber	roots/end of season after foliage dies	Root similar to potato with rough skin. Skin. Use like potato. Also called blue taro. Can be confused with Taro (<i>Colocasia esculenta</i>) which is also used in Chinese cuisine.
Water Chestnuts Boh chi <i>Eleocharis dulcis</i>	7 diam 5 deep	1	3-5	after danger of frost	tubers or corms	"nuts" (corms)/ after 6 months of growth	"Chestnuts" form in mud at base of stem. Walnut size or smaller. Peel tough skin, eat raw or cooked.

For more information on the subject discussed in this publication, consult your local office of the Purdue University Cooperative Extension Service.

It is the policy of the Purdue University Cooperative Extension Service, David C. Petritz, Director, that all persons shall have equal opportunity and access to programs and facilities without regard to race, color, sex, religion, national origin, age, marital status, parental status, sexual orientation, or disability. Purdue University is an Affirmative Action employer. This material may be available in alternative formats. <http://www.agcom.purdue.edu/AgCom/Pubs/menu.htm>